

Functional movement test scores improve following a standardized off-season intervention program in professional football players

Q1 K. Kiesel^{1,2}, P. Plisky², R. Butler¹

¹University of Evansville, Evansville, Indiana, USA, ²ProRehab PC, Evansville, Indiana, USA

Corresponding author: Kyle Kiesel, PT, PhD, ATC CSCS, University of Evansville, Wallace Graves Hall 206, Evansville, Indiana 47722, USA. Tel: +812 479 2646, Fax: +812 479 2717, E-mail: kk70@evansville.edu

Accepted for publication 30 July 2009

The purpose of this study was to determine if an off-season intervention program was effective in improving Functional Movement Screen™ (FMS) scores in professional American football players. Pre- and post-intervention FMS scores were obtained on 62 subjects who completed a 7-week off-season intervention program. A repeated measures ANOVA was conducted to determine the effectiveness of the training program on FMS scores. A chi-square was performed to determine if there were a greater number of players who met the injury threshold and if asymmetries were reduced following intervention. Logistic regression was used to

predict what factors were associated with failure (post-test score of <14). There was a positive main effect for time ($P < 0.01$) and a greater number of individuals with a score >14 following than intervention. At post-test, 41 players were free of asymmetry as compared with 31 at the pre-test. The strongest predictor of program failure was a low squat score at pre-test. This study demonstrated that fundamental movement characteristics do change with a standardized intervention. Further research is required to determine if injury risk is reduced when a player's score improves beyond the established cut-off of 14 and/or asymmetry is resolved.

There are limited published reports on injury risk factors for professional football players (Powell, 1987). The most prominent risk factor identified in prospective injury risk studies in athletics is previous injury (Turbeville et al., 2003; Tyler et al., 2006). Previous injury is a complex risk factor implying that something fundamentally changes following injury. It is possible that following injury there is a fundamental change in motor control that is not measurable with a single traditional impairment measurement such as strength or range of motion. In an attempt to capture the complex construct of motor control, movement oriented tests that assess functional movement and dynamic balance have been developed (Cook et al., 2006; Plisky et al., 2006). Tasks requiring coordinated full body movements may, in part, help us understand what fundamentally changes following injury. One test that has been established to assess neuromuscular impairments with movement is the Functional Movement Screen™ (FMS).

The FMS is a test of seven fundamental movements that has previously been determined to be able to predict injury in professional football players. Previous research has shown that players scoring lower than a 14 on the FMS were more likely to be injured than those scoring above 14 (Kiesel et al., 2007). A follow-up study to this initial investigation, validated the previous finding and in addition observed that players with any asymmetry on the FMS, regardless of total

score, had $2.3 \times$ greater risk for injury (Kiesel et al., 2008). While it appears the FMS may be beneficial in injury prediction, no studies have assessed whether changes in an individual's FMS score can be achieved with a training program.

As a result of the aforementioned literature, the purpose of this study was to determine if an off-season intervention program is effective in improving FMS scores in professional American football players. In addition, this study wanted to examine if there was a greater percentage of players above the injury threshold of a score of 14 at the end of the study compared with the beginning of the study. Because, right and left asymmetry in the FMS has also been related to injury we also examined if more players were free of asymmetry at the end of the study compared with the beginning of the study. Finally, we wanted to determine if it was possible to predict who would not improve their score above the injury threshold score using data from the pre-screen.

Methods

This study was completed on a sample of 62 healthy American football players from one professional team who participated in a structured off-season strength and conditioning program. In addition to traditional strength and conditioning components, the program also included an individualized interven-

	S M S	1 0 3 8	B	Dispatch: 6.10.09	Journal: SMS	CE: Sangeeta
	Journal Name	Manuscript No.		Author Received:	No. of pages: 6	PE: Chris/Elan

tion program which was prescribed based on the individual's score on the FMS™ as described by Cook (2001).

FMS

The FMS is a comprehensive screen that assesses quality of fundamental movement patterns to identify an individual's limitations or asymmetries (Cook et al., 2006). The FMS includes seven tests which are scored on a 0–3 ordinal scale. The seven tests of the FMS are the squat, hurdle step, lunge, shoulder mobility, active straight leg raise, push up and rotary stability. A score of 3 indicates the movement was completed as instructed and is free of compensation and pain. A score of 2 indicates the subject could complete the movement pain-free but with some level of compensation, while a score of 1 indicates the subject could not complete the movement as instructed. A score of 0 is assigned if the subject experiences pain with any portion of the movement. The reliability of the FMS has been established in a recent study (Minick et al., 2009) which reported weighted kappa values to range from 0.80 to 1.00.

Of the seven tests that comprise the FMS, five of them (hurdle step, lunge, shoulder mobility, active straight leg raise and rotary stability) are performed and scored separately for the right and left sides of the body. Because of the established relationship between neuromuscular asymmetry (right vs left differences) and injury risk, it is common practice to track right vs left sided differences on the FMS. Asymmetry is dichotomized for research purposes and if an asymmetry exists in an individual's score it is prioritized in the corrective program in an attempt to achieve the goal of symmetrical movement on all five of the right/left FMS tests.

Off-season training program

After initial FMS scores were obtained, subjects were prescribed an individual training program, based on their FMS score, designed to correct the identified movement deficits. This program consisted of a "movement preparation" component that included self and partner stretching and self-administered trigger point treatment done using The Stick (The Stick/RPI Atlanta, Georgia, USA) of major muscle groups thought to be contributing to the dysfunctional movement patterns. Individual "corrective exercises" were also prescribed and performed with the supervision and instruction of the strength and conditioning staff as needed. The corrective exercises are intended to utilize the range of motion gained from the stretching and allow for motor learning to occur as the subject moves through previously limited patterns of movement. Additionally, the corrective exercises are designed to stimulate natural core muscle activation to enhance the relationship between core muscle function and fundamental movement (Cook & Fields, 1997).

The goal was for the subjects to become independent with performing both components of the program. The exercises were progressed in a standardized manner with the goal of achieving the best possible movement strategy related to each of the seven movements tested. The program also included traditional strength and conditioning activities. See Table 1 for an example of an individual intervention program utilized in this study that incorporates movement preparation and corrective exercise as defined.

Athletes were required to participate in the 7-week program which included supervised sessions 4 days/week and two optional sessions per week. The program followed a structured progression with the goal of completing the final phase of corrective exercises as part of the routine strength and conditioning program. Post-test FMS scores were obtained at the completion of the program.

Statistical analysis

For the purposes of analysis, the subjects were grouped by position with lineman and linebackers (lineman, $n = 32$) in one group and all others position players in another group (non-lineman, $n = 30$). Mean and standard deviations for the pre-test and post-test FMS scores were calculated for each group. To determine if there was a change in FMS scores following the intervention period, a repeated measures ANOVA with the significance level set at $P < 0.05$ was conducted with the repeated measure being time (pre-test and post-test scores) and the between subjects variable being group (lineman and non-lineman). A one-way chi-square was performed to determine if there was a significant increase in the number of players scoring above the injury threshold score of 14 following the program. A similar analysis was performed to determine if there was an increase in the number of players that were free from asymmetry at the end of the program. For the chi-square analysis, the expected distribution utilized for the analysis was the distribution that was obtained at baseline. Finally, a logistic regression was utilized, using the variables position, asymmetry present and baseline score on the deep squat in order to predict which individuals did not improve their score to above the injury score threshold at the end of the program. Further rationale for the selection of these variables is highlighted in the discussion section of the manuscript.

Results

There was no time by position interaction for the FMS scores over the course of the training program (Fig. 1, $F = 0.08$, $P = 0.78$), however there was a main effect of time (Fig. 1, $F = 180.4$, $P < 0.01$). The mean pre-test score for the lineman was 11.8 (SD 1.8) and for the non-lineman group was 13.3 (SD 1.9). The mean post-test score for the lineman was 14.8 (SD 2.4) and for the non-lineman group was 16.3 (SD 2.4). Combined data for all subjects for the pre-test and post-test for each component of the FMS are presented in Fig. 2.

At the end of the intervention, a greater number of players exhibited a score that improved to above the injury threshold compared with before the intervention ($\chi^2 = 164.9$, $P < 0.01$). Thirty-nine subjects exhibited a score that exceeded the injury risk threshold at the end of the program compared with seven that exhibited a score > 14 at baseline (Table 2).

In regards to the changes in symmetry scores, a greater percentage of players were free from asymmetry at the end of the study as compared with the beginning of the study ($\chi^2 = 7.8$, $P = 0.01$). Following the intervention, 42 players were free of asymmetry compared with the 31 players that demonstrated at least one asymmetry at baseline (Table 2).

The results of the logistic regression suggest that failure to improve above the injury threshold score of 14 can be predicted by a low deep squat score during the initial data collection. Neither position nor the presence of asymmetry was a significant predictor of whether or not the players score was able to be improved above the score of 14.

Q8 Table 1. Example of an individual off-season intervention program based on pre-test Functional Movement Screen™ score

Test	Score	Final	Movement prep	Corrective exercise
Deep squat	2	2		
Hurdle step	R-2 L-3	2	 <p>Anterior thigh stick work</p>	 <p>Straight leg bridge</p>
In-line lunge	R-2 L-2	2	 <p>Stride Self-Stretch</p>	 <p>Single leg stance w/core engagement</p>
Shoulder mobility	R-3 L-3	3		
Active straight leg raise	R-1 L-2	1	Anterior thigh Stick work (see above)	
			 <p>Straight leg partner stretch</p>	 <p>Leg lowering w/core engagement</p>
			 <p>Leg lowering progression</p>	 <p>Single-leg toe touch w/stick</p>
Trunk stability push-up	3	3		
Rotary stability	R-2 L-2	2		
Composite		15		
Asymmetry?		Yes		

Discussion

There is currently no evidence of the ability to change scores on the FMS based on an off-season training program. The results of this study support that an off-season training program can significantly improve scores on the FMS. In addition, the off-season

training program significantly increased the percentage of players who scored above the previously determined injury threshold score and also significantly improved the percentage of players who were free of asymmetry. Finally, the results of the study suggest that one of the primary predictors of the inability to improve a player's score above the injury

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000




Fig. 1. Composite Functional Movement Screen™ (FMS) scores for non-lineman and lineman before and after the 7-week intervention.




Fig. 2. Mean pre/post-values for the all subjects for each individual component of the Functional Movement Screen™ (FMS). DS, deep squat; HS, hurdle step; ILL, in-line lunge; ASLR, active straight leg raise; SM, shoulder mobility; TSPU, trunk stability push-up; RS, rotary stability.

Table 2. Changes in number of players composite Functional Movement Screen™ score (FMS) and presence of asymmetries from pre-to post-test

	Pre-test	Post-test	χ^2	P
FMS ≥ 14	7	30	164.9	<0.01
14 > FMS	55	32		
Asymmetry	31	20	7.8	0.01
No asymmetry	31	42		

threshold score was a low score on the deep squat during the pre-test.

While there was no position by time interaction, all players, on average, exhibited an 11% increase in their FMS score as a result of the off-season training program. This suggests that functional movement, as measured by the FMS, can be improved when a

structured intervention is applied. While this result is promising, caution should be taken when interpreting this result as no control group was included in this study. It is possible that the changes observed in this study were an affect of repeated measures of test taken at the start and again at the completion of the program. However, while there was no control group, the change in FMS scores from pre-test to post-test ranged from -1 to $+7$ suggesting not all players systematically improved their score as a function of repeated measures.

It is difficult to directly compare our results to other research as we are unaware of any other published study to date which has tested the FMS in this manner. Our results are similar to those reported using other body relative movement tests. Significant improvement in the Star Excursion Balance Test has been reported following standardized intervention programs including Pilates, (English & Howe, 2007) dynamic hop and balance training (McKeon et al., 2008) and functional rehabilitation programs (Hale et al., 2007).

Potentially more important than the absolute improvement in scores was the significant increase in the number of players who scored above the injury threshold at the post-test. At the pre-test only 11% of the players exhibited a score >14 while 63% of players exhibited a score >14 at the post-test. This finding may be of benefit to sports medicine professional who work with professional football players as 52% of the players in the program were able to improve their score from below to above the established threshold score for injury risk (≤ 14).

Although a significant number of players improved their score to >14 , the question remains as to why 20 subjects failed to improve their score enough to exceed the threshold of 14? In order to address this issue, a logistic regression model was developed in an attempt to identify if any factors were present which could predict failure (not improving to above a score of 14). The sample size of 32 allowed us to only include three factors into the model which were selected by the research group to include position, presence of an asymmetry and scoring a 1 on the Deep Squat test. Position was chosen because of the different movement demands inherent to the different positions in this sport. Asymmetry was chosen because it is a well-known injury risk factor and because we hypothesized that the presence of an asymmetry may be a contributing factor to overall movement limitations. The squat test was included because it involves a balance of mobility (ankles, knees, hip, thoracic spine and shoulders) and stability (pelvic stability to allow the hips to break parallel while maintaining an upright trunk). As a result of the complexity of this task, failure of this test may be an indication of a substantial movement dysfunction in need of an individual examination

rather than a basic screen and exercise program. The results indicated that those players scoring a 1 on the Deep Squat Test were approximately 5 × more likely to fail (OR 5.27, 95% CI 1.49–18.65) while the other factors were not shown to be significant predictors of failure. The fact that not all subjects improved their score beyond the threshold of 14 (failure) and that poor deep squat performance was related to failure, suggests that a more robust program may be needed for these individuals to normalize their movement patterns. Perhaps an individual assessment may be required to identify the underlying neuromuscular factors contributing to the persistent dysfunctional movement.

The final area that was examined was asymmetry of movement, due to the fact that musculo-skeletal asymmetry is a well established risk factor for injury (Ekstrand & Gillquist, 1983; Knapik et al., 1991; Baumhauer et al., 1995; Nadler et al., 2000; Soderman et al., 2001; Plisky et al., 2006; Myer et al., 2008; Yeung et al., 2009). At the pre-test 50% of the players demonstrated asymmetrical movement and this was reduced to 32% following the interventions. This result would suggest that the off-season program was not only able to normalize for poor total body movements but was also able to neutralize some of the asymmetries that were measured in the players.

Some additional limitations of this study should be noted. There was a lack of a control group, therefore, no clear cause/effect relationship can be established without randomization and control. A second potential limitation is associated with the repeated measurement testing which has been previously discussed. Additionally, there was not control for co-interventions. It is possible that some of the subjects were receiving additional intervention such as personal massage or chiropractic care that may have influenced the outcome of the study. Finally, there was no short-term follow-up, perhaps the changes in scores observed occurred much earlier in the program than at the terminal 7-week point. Additional research should focus on examining the change in the FMS as it relates to dosage of a corrective exercise program.

Although the specific results of this study cannot be generalized beyond the population tested, limitations and asymmetries in fundamental movement patterns which are measured by the FMS are basic components of human movement and are not sport

specific. Therefore, it may be beneficial to perform additional studies in other active populations.

In conclusion, this study demonstrates that movement can improve with an individualized exercise program. Injury prevention and performance enhancement programs should consider including the FMS or a similar movement screen and associated exercises to normalize dysfunctional movement with the goal of injury reduction and performance improvement.

Perspectives

It is known that previous injury is the most robust risk factor in sport. Although individual components of athletic movement such as strength, flexibility and power are routinely addressed in contemporary rehabilitation, this may not serve as a protective factor against injury. Motor control changes exist following injury and movement oriented testing and training programs to address these deficits are needed in sports (Minick et al., 2009). Recent research (Paterno, 2009), reported that there was no relationship between strength, power, or flexibility and performance on the star excursion balance test. This may indicate that movement oriented tests measure a separate construct which could be helpful in understanding more about motor control changes following injury. The FMS has identified individuals at risk of injury in professional football but it has yet to be determined if an individual's FMS score could be altered by a training program. The results of this study suggest that American professional football players can improve their score with an off-season training program. It has yet to be determined if these improvements will occur in other active populations and if prospective improvements in the FMS actually reduce injury risk. The FMS may be a tool to assist the sports medicine professional in reducing injury risk.

Key words: assessment, injury prevention, motor control.

Acknowledgements

The authors would like to thank Jeff Fish, Director of Athletic Performance and Bill Hughan Assistant Strength Coach both of the Atlanta Falcons for their assistance with data collection for this manuscript.

References

- | | | |
|---|---|---|
| Baumhauer JF, Alosa DM, Renstrom AF, Trevino S, Beynonn B. A prospective study of ankle injury risk factors. <i>Am J Sports Med</i> 1995; 23(5): 564–570. | Cook G. Athletic body balance. Champaign, IL: Human Kinetics, 2001. | assessment of function – part 1. <i>NAJSPT</i> 2006; 1: 62–72. |
| | Cook G, Burton L, Hogenboom B. The use of fundamental movements as an | Cook G, Fields K. Functional training for the torso. <i>Strength Cond</i> 1997; 19(2): 14–19. |

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

- Ekstrand J, Gillquist J. Soccer injuries and their mechanisms: a prospective study. *Med Sci Sports Exerc* 1983; 15(3): 267–270.
- English T, Howe K. The effect of pilates exercise on trunk and postural stability and throwing velocity in college baseball pitchers: single subject design. *NAJSPT* 2007; 2: 8–19. **Q3**
- Hale SA, Hertel J, Olmsted-Kramer LC. The effect of a 4-week comprehensive rehabilitation program on postural control and lower extremity function in individuals with chronic ankle instability. *J Orthop Sports Phys Ther* 2007; 37: 303–311.
- Kiesel K, Plisky P, Kersey P. Functional movement test score as a predictor of time-loss during a professional football team's pre-season American college of sports medicine annual conference. Indianapolis, IN, 2008. **Q4**
- Kiesel K, Plisky P, Voight M. Can serious injury in professional football be predicted by a preseason functional movement screen? *North Am J Sports Phys Ther* 2007; 2: 147–158.
- Knapik JJ, Bauman CL, Jones BH, Harris JM, Vaughan L. Preseason strength and flexibility imbalances associated with athletic injuries in female collegiate athletes. *Am J Sports Med* 1991; 19(1): 76–81.
- McKeon PO, Ingersoll CD, Kerrigan DC, Saliba E, Bennett BC, Hertel J. Balance training improves function and postural control in those with chronic ankle instability. *Med Sci Sports Exerc* 2008; 40: 1810–1819.
- Minick K, Burton L, Butler R, Kiesel K. A reliability study of the functional movement screen. *Natl J Strength Cond Res* 2009, in press.
- Myer GD, Ford KR, Paterno MV, Nick TG, Hewett TE. The effects of generalized joint laxity on risk of anterior cruciate ligament injury in young female athletes. *Am J Sports Med* 2008; 36(6): 1073–1080.
- Nadler SF, Malanga GA, DePrince M, Stitik TP, Feinberg JH. The relationship between lower extremity injury, low back pain, and hip muscle strength in male and female collegiate athletes. *Clin J Sport Med* 2000; 10(2): 89–97.
- Paterno MVSL, Ford K, Hewett T. Contribution of lower extremity strength and postural stability to performance on the star excursion balance test combined section conference of the American physical therapy association. Las Vegas, NV, 2009.
- Plisky PJ, Rauh MJ, Kaminski TW, Underwood FB. Star excursion balance test as a predictor of lower extremity injury in high school basketball players. *J Orthop Sports Phys Ther* 2006; 36: 911–919.
- Powell J. Incidence of injury associated with playing surfaces in the national football league. *J Athl Train* 1987; 22: 202–206.
- Soderman K, Alfredson H, Pietila T, Werner S. Risk factors for leg injuries in female soccer players: a prospective investigation during one out-door season. *Knee Surg Sports Traumatol Arthrosc* 2001; 9(5): 313–321. **Q5**
- Turbeville S, Cowan L, Owen W. Risk factors for injury in high school football players. *Am J Sports Med* 2003; 31: 974–980.
- Tyler TF, McHugh MP, Mirabella MR, Mullaney MJ, Nicholas SJ. Risk factors for noncontact ankle sprains in high school football players: the role of previous ankle sprains and body mass index. *Am J Sports Med* 2006; 34: 471–475.
- Yeung SS, Suen AM, Yeung EW. A prospective cohort study of hamstring injuries in competitive sprinters: preseason muscle imbalance as a possible risk factor. *Br J Sports Med* 2009. **Q7**
- Q6**

10/06/2009 2:02:20 PM
 10/06/2009 02:02:15 -00219
 Byres:6 PAGES:6
 operator=jhm.Christine

